

Longer heat waves tend to have broad footprints

Day+1

50°N

40°N

30°N


120°W

100°W


80°W

Composite
Tmax and
700mb height
Anomalies from
31 historical
3d and longer
heat wave
events
(1948-2005)


Composite 700ht Anoms, 31 Hi Tmax dates lasting >= 3days (days when >= 9 Sasha stns have hot days by 99%ile criteria)




#### California Heat Waves: 2006 vs. historical record




Locally extreme temperature exceeding a high percentile threshold t\* (e.g. 99%-ile)






CA heat wave index: %-age of stations under extreme heat during day and night



Concentration of heat wave events in June-September


Composite 700ht Anoms, 31 Hi Tmax dates lasting >= 3days (days when >= 9 Sasha stns have hot days by 99%ile criteria)




Circulation during
A heat wave episode

700 mb height anomalies composited 4 days before through 6 days after start date


#### Comp Dly Stn Tmax DM, 31 Hi Tmax dates lasting >= 3 days





# West-wide expression: Tmax anomalies 3d before through 3d after




July 2006
Temperature
Humidity
Circulation
anomalies









## Near-surface humidity anomalies during July 2006



Fresno, CA (KFAT) No. of Tdew/Tair obs (3PM PST)





#### San Diego, CA (KSAN) No. of Tdew/Tair obs (3PM PST)





# Daytime and Nighttime Statewide Heatwave Activity

Daytime Maximum Temperature events: episodic, not prominent every year Nighttime Minimum Temperature events: have increased over last 50 years, 2006 had remarkably warm Tmins











Medium range forecast skill 15d-1d in advance


Los
Angeles
area,
Heat spell
(3days and
longer)
days

Dave load

### Climate Change Projections Heat wave days

















- Locally extreme temperature exceeding a high percentile threshold t\* (e.g. 99%-ile)
- Local heat wave magnitude and duration reflected in summing exceedance over threshold: ∑(t<sub>i</sub> – t\*), for all days (i) in a season such that (t<sub>i</sub> > t\*) in a season
- Spatial extent of locally extreme conditions

Seasonal view of daily extremes in time and space:

individual seasons vs. multiyear climatology

 Spatial extent (weighted by population) of locally extreme conditions is computed for each year as percentage of stations at which exceedance sums are above the
 climatological threshold

% of stations

1950

1960


1970

1980

1990

2000

99<sup>th</sup> %-ile of climatological exceedance sums





This gives the percentage of stations where heatwave activity was extreme

## **Tmax**

 Locally extreme temperature exceeding a high percentile threshold t\* (e.g. 99%-ile)

Time series view







Bars: 1948-2005 Tmax histogram, Circles: 2006 Tmax histogram, and Vertical line: all-yr 99<sup>th</sup> %-ile

Each season is compared to local climatology

Local heat wave
 magnitude and duration
 reflected in summing
 exceedances over
 threshold: ∑(t<sub>i</sub> – t\*), for all
 days (i) in a season such
 that (t<sub>i</sub>T\*nt\*) series view



99<sup>th</sup> %-ile of climatological exceedance sums




Exceedance sums are computed for each year at each station on our network that is weighted towards most populated areas


## **Tmin**

» Locally extreme temperature exceeding a high percentile threshold t\* (e.g. 99%-ile)

#### Time series view



#### Sacramento Tmax



Bars: 1948-2005 Tmax histogram, Circles: 2006 Tmin histogram, and Vertical line: all-yr 99th %-ile

Each season is compared to local climatology

### Composite SST Anoms (deg F), Hi Tmax Dates >= 3 days

